

A diffuse-domain-based numerical method for a chemotaxis-fluid model

Chenxi Wang

*Department of Mathematics, Harbin Institute of Technology,
Harbin 150001, P. R. China*

*Department of Mathematics,
Southern University of Science and Technology,
Shenzhen 518055, P. R. China
wangcx2017@mail.sustech.edu.cn*

Alina Chertock

*Department of Mathematics, North Carolina State University,
Raleigh, NC 27695, USA
chertock@math.ncsu.edu*

Shumo Cui^{*,§}, Alexander Kurganov^{†,¶,**} and Zhen Zhang^{‡,||}

**Department of Mathematics, International Center for Mathematics,
Southern University of Science and Technology,
Shenzhen 518055, P. R. China*

*†Department of Mathematics, International Center for Mathematics,
Guangdong Provincial Key Laboratory of Computational Science
and Material Design, Southern University of Science and Technology,
Shenzhen 518055, P. R. China*

*‡Department of Mathematics,
Guangdong Provincial Key Laboratory of Computational Science
and Material Design, National Center for Applied Mathematics (Shenzhen),
Southern University of Science and Technology,
Shenzhen 518055, P. R. China*

§cuism@sustech.edu.cn

¶alexander@sustech.edu.cn

||zhangz@sustech.edu.cn

Received 16 September 2022

Revised 28 November 2022

Accepted 14 December 2022

Published 23 February 2023

Communicated by N. Bellomo

In this paper, we consider a coupled chemotaxis-fluid system that models self-organized collective behavior of oxytactic bacteria in a sessile drop. This model describes the biological chemotaxis phenomenon in the fluid environment and couples a convective

**Corresponding author.

chemotaxis system for the oxygen-consuming and oxytactic bacteria with the incompressible Navier–Stokes equations subject to a gravitational force, which is proportional to the relative surplus of the cell density compared to the water density.

We develop a new positivity preserving and high-resolution method for the studied chemotaxis-fluid system. Our method is based on the diffuse-domain approach, which we use to derive a new chemotaxis-fluid diffuse-domain (cf-DD) model for simulating bioconvection in complex geometries. The drop domain is imbedded into a larger rectangular domain, and the original boundary is replaced by a diffuse interface with finite thickness. The original chemotaxis-fluid system is reformulated on the larger domain with additional source terms that approximate the boundary conditions on the physical interface. We show that the cf-DD model converges to the chemotaxis-fluid model asymptotically as the width of the diffuse interface shrinks to zero. We numerically solve the resulting cf-DD system by a second-order hybrid finite-volume finite-difference method and demonstrate the performance of the proposed approach on a number of numerical experiments that showcase several interesting chemotactic phenomena in sessile drops of different shapes, where the bacterial patterns depend on the droplet geometries.

Keywords: Chemotaxis; Navier–Stokes equations; bioconvection; diffuse-domain approach; finite-volume method; finite-difference method.

AMS Subject Classification: 65M85, 65M06, 65M08, 92C17, 76Z99

1. Introduction

In this paper, we study the following coupled chemotaxis-fluid system in a sessile drop⁴⁴:

$$\begin{aligned} n_t + \mathbf{u} \cdot \nabla n + \chi \nabla \cdot [nr(c)\nabla c] &= D_n \Delta n, & c_t + \mathbf{u} \cdot \nabla c &= D_c \Delta c - n\kappa r(c), \\ \rho(\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u}) + \nabla p &= \eta \Delta \mathbf{u} - n \nabla \Phi, & \nabla \cdot \mathbf{u} &= 0, \end{aligned} \quad (1.1)$$

where n and c are the concentrations of bacteria and oxygen, respectively, κ is the oxygen consumption rate, and $\mathbf{u} = (u, v)^\top$ is the velocity field of a fluid flow governed by the incompressible Navier–Stokes equations with the density ρ , pressure p , and viscosity η . In the fluid equation, $\nabla \Phi := V_b g(\rho_b - \rho)\mathbf{z}$ describes the gravitational force exerted by a bacterium onto the fluid along the upwards unit vector \mathbf{z} proportional to the volume of the bacterium V_b , the gravitation acceleration $g = 9.8 \text{ m/s}^{-2}$, and the density of bacteria is ρ_b (bacteria are about 10% denser than water).

In Eq. (1.1), both the bacteria and oxygen are convected by the fluid and diffuse with their respective diffusion coefficients D_n and D_c . The bacteria are active as long as a sufficient oxygen supply is available: this is measured by a dimensionless cut-off function $r(c)$, which can be modeled, for instance, by

$$r(c) = \begin{cases} 1, & c \geq c^*, \\ 0, & c < c^*, \end{cases} \quad (1.2)$$

where c^* is an inactivity threshold. The active bacteria both consume the oxygen and, in a chemotactic response, are directed towards a higher oxygen concentration with the rate proportional to the chemotactic sensitivity χ .




Fig. 1. A sketch of the sessile drop domain Ω .

A typical shape of the sessile drop is depicted in Fig. 1. We stress that the boundary conditions on n , c , and \mathbf{u} are essential to ensure that the solutions of (1.1) match well the experimental observations. We shall consider the following mixed boundary conditions: the boundary conditions on the top interface Γ describe the fluid-air surface, which is stress-free, allows no cell flux, and has saturated air oxygen concentration c_{air} :

$$\begin{aligned} \boldsymbol{\nu} \cdot \mathbf{u} &= 0, \quad \boldsymbol{\nu} \cdot \nabla(\mathbf{u} \cdot \boldsymbol{\tau}) = 0, \\ (\chi n \nabla c - D_n \nabla n) \cdot \boldsymbol{\nu} &= 0, \quad c = c_{\text{air}}, \quad \forall (x, y) \in \Gamma, \end{aligned} \quad (1.3)$$

where $\boldsymbol{\nu}$ and $\boldsymbol{\tau}$ are the unit outward normal and tangential vectors on Γ . A no-slip boundary condition is applied on the bottom surface ($\partial\Omega_{\text{bot}}$) and there is no flux of cells or oxygen through $\partial\Omega_{\text{bot}}$:

$$\mathbf{u} = \mathbf{0}, \quad \nabla n \cdot \boldsymbol{\nu} = 0, \quad \nabla c \cdot \boldsymbol{\nu} = 0, \quad \forall (x, y) \in \partial\Omega_{\text{bot}}. \quad (1.4)$$

The chemotaxis-fluid system (1.1) belongs to a class of cross-diffusion models in complex environment. A classical cross-diffusion model is the Patlak–Keller–Segel (PKS) system,^{32, 33, 38} which paved the foundation for the mathematical modeling of chemotaxis. In the following decades, specializations and variations of the PKS system have been developed (see the reviews papers^{5, 24} and references therein) together with the coupled chemotaxis-fluid models.^{17, 23, 25, 36, 44, 46, 47} The aforementioned chemotaxis models were usually obtained by means of heuristic interpretation of macroscopic chemotaxis phenomena. Recently, the framework of micro-macro derivation has been applied to chemotaxis and chemotaxis-fluid models.^{6, 8} The micro-macro framework unified the derivation of chemotaxis models with other models describing collective dynamics of smaller-scale entities, for example, crowd dynamics,⁷ virus pandemic,⁹ etc.

There is a large body of research works investigating the qualitative properties (for example, the global existence, boundedness, stability properties of equilibrium configurations, and blow-up solutions) of the PKS system and its variations; see, e.g. Ref. 27 and references therein. Recently, stability analysis and dynamics of the chemotaxis-fluid system (1.1) with a deformed free-surface in a shallow chamber were studied in Ref. 14. In particular, a detailed linear stability analysis of a steady-state cell and oxygen concentration distribution was performed there. A global weak solution of (1.1) can be obtained as a limit of smooth solutions of a regularized chemotaxis-fluid system under suitable boundary and initial conditions; see Ref. 31. Moreover, it has been recently proven in Ref. 11 that in the one- or two-dimensional

cases, the system (1.1) has a unique global classical solution provided $r(c)$ is constant and the following boundary conditions: no-slip for u , no-flux for n , and Robin for c , are used in a smoothly bounded convex domain. In the three-dimensional case, the existence of a global weak solution in a drop shaped domain has been shown and a uniform in time energy bound has been established. The link between the diffusion and the cross-diffusion terms in the chemotaxis equation would also affect the analytical results for more general chemotaxis-fluid systems.³¹ For recent analytical results related to (1.1)–(1.4) or general cross-diffusion models, we refer the reader to Refs. 8 and 31 and references therein.

Qualitative properties of chemotaxis-fluid models were also investigated numerically. In Ref. 15, the system (1.1) was studied in a simplified, rectangular shaped domain subject to the same top and bottom boundary conditions as in (1.3) and (1.4), respectively, and periodic boundary conditions in the horizontal direction. Several phenomena of sinking, merging and stationary plumes were discovered in Ref. 15 by numerically solving (1.1) using a high-resolution hybrid finite-volume finite-difference method. In Ref. 16, an upwind finite-element method was developed and used to investigate the pattern formation and the hydrodynamical stability of the system (1.1) for the same simplified setup. In Ref. 28, a fully decoupled, linear and positivity preserving finite-element method for solving the chemotaxis-Stokes equations has been recently developed for a similar setup. In Ref. 18, the chemotaxis-fluid model without the discontinuous oxygen cut-off function $r(c)$ has been considered, for which a finite-element method has been constructed, optimal error estimates have been established, and convergence towards regular solutions has been proved. In Refs. 29 and 30, a generalized chemotaxis–diffusion–convection model, which includes the dynamic free surface and appropriate boundary conditions, has been proposed together with a numerical method, which uses a time-dependent grid and incorporates surface tension and a dynamic contact line. However, extensive studies of bacterial swimming and oxygen transport dynamics in more complex geometries are still in great demand, as they can shed light on qualitative analysis.

The main goal of this paper is to develop a robust and accurate numerical method for the chemotaxis-fluid system in the sessile drop domain. To this end, we extend a diffuse-domain approach to the system (1.1) and construct a new chemotaxis-fluid diffuse-domain (cf-DD) model, which we numerically solve using a second-order hybrid finite-volume finite-difference method.

The diffuse-domain method was proposed in Ref. 35 following the idea of the smoothed boundary method previously introduced in Refs. 12 and 13 as a powerful numerical tool for solving diffusion equations with no-flux boundary conditions imposed at irregular boundaries within the computational domain. The diffuse-domain method can be applied to a variety of PDEs in both stationary and moving complex geometries with Dirichlet, Neumann, or Robin boundary conditions. The key idea of the method is to place the complex geometry into a larger rectangular domain, introduce a smoothed characteristic function of the original domain,

and reformulate the original PDE(s) on the extended domain with the help of additional source terms, which reflect the contribution of the original boundary conditions. It has been shown in Refs. 22, 34, 35, 42, 49 and 50 that the reformulated diffuse-domain model asymptotically converges to the original PDE(s) as the thickness of the diffuse-domain interface tends to zero. The main advantage of the diffuse-domain method is that the reformulated model can be solved using standard numerical methods even for very complex domains (with moving boundaries). For example, the diffuse-domain method has been successfully applied to several quite sophisticated two-phase flow models; see Refs. 2, 3, 43 and 48.

We first follow the diffuse-domain approach and derive a cf-DD model, for which we perform an asymptotic analysis and show that it converges to the original chemotaxis-fluid model as the thickness of the diffuse-domain interface shrinks to zero. We then use the proposed cf-DD model to simulate bioconvection in complex droplet geometries using a numerical method, which is derived as follows. The modified cell density equation is numerically solved by a semi-discrete second-order finite-volume upwind method (introduced in Ref. 15) combined with a second-order strong stability-preserving multistep ODE solver, which can be found in, e.g. Ref. 19. The resulting fully discrete scheme is shown to preserve the positivity of cell density. The modified Navier–Stokes and oxygen concentration equations are discretized using a second-order projection finite-difference method, combined with the second-order BDF-like method for the time evolution. The proposed numerical method produces results which, in the middle part of the considered droplets, qualitatively similar to those reported in Ref. 15. Using the new method, we were able to capture complicated dynamics of the bacteria cells including emergence of plumes and their evolution in complex droplet geometries.

The rest of the paper is organized as follows. In Sec. 2, we describe a non-dimensional version of the coupled chemotaxis-fluid system (1.1) and introduce typical values of the scaling parameters to be used in our numerical simulations. In Sec. 3, we present the reformulated cf-DD model. In Sec. 4, we introduce the numerical method for the cf-DD system and discuss its implementation. In Sec. 5, we report several numerical experiments illustrating the performance of the proposed diffuse-domain-based numerical method. Finally, in Sec. 6, we give few concluding remarks and discuss perspectives of our future work.

2. Scaling and Setup

We denote by L a characteristic length (we may choose, for instance, L to be the maximum height of the drop; see Fig. 1) and the characteristic cell density by n_r . Rescaling the variables as in Refs. 15 and 44

$$\begin{aligned} \mathbf{x}' &= \frac{\mathbf{x}}{L}, & t' &= \frac{D_n}{L^2} t, & c' &= \frac{c}{c_{\text{air}}}, \\ n' &= \frac{n}{n_r}, & p' &= \frac{L^2}{\eta D_n} p, & \mathbf{u}' &= \frac{L}{D_n} \mathbf{u}, \end{aligned} \tag{2.1}$$

leads to the five dimensionless parameters α , β , γ , δ , and the Schmidt number Sc :

$$\begin{aligned}\alpha &:= \frac{\chi c_{\text{air}}}{D_n}, \quad \beta := \frac{\kappa n_r L^2}{c_{\text{air}} D_n}, \\ \gamma &:= \frac{V_b n_r g(\rho_b - \rho) L^3}{\eta D_n}, \quad \delta := \frac{D_c}{D_n}, \quad \text{Sc} := \frac{\eta}{D_n \rho},\end{aligned}\tag{2.2}$$

which characterize the system (1.1)–(1.4). Three of the parameters in (2.2), namely, α , δ , and Sc , are determined by the properties of bacteria, fluid and air. Typical values for *Bacillus subtilis* in water are $\alpha = 10$, $\delta = 5$, and $\text{Sc} = 500$; see, e.g. Ref. 44. The remaining two parameters β and γ depend also on the chosen length scale L and the reference cell density n_r , and thus will be varied in the numerical examples reported in Sec. 5.

Dropping the primes from the dimensionless quantities in (2.1) yields the following non-dimensional version of the governing chemotaxis-fluid system:

$$n_t + \nabla \cdot (\mathbf{u}n) + \alpha \nabla \cdot [r(c)n \nabla c] = \Delta n,\tag{2.3}$$

$$c_t + \mathbf{u} \cdot \nabla c = \delta \Delta c - \beta r(c)n,\tag{2.4}$$

$$\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u} + \text{Sc} \nabla p = \text{Sc} \Delta \mathbf{u} - \text{Sc} \gamma n \mathbf{z},\tag{2.5}$$

$$\nabla \cdot \mathbf{u} = 0.\tag{2.6}$$

This system is considered on a sessile drop domain Ω subject to the initial data

$$n(x, y, 0) = n_0(x, y), \quad c(x, y, 0) = c_0(x, y), \quad \mathbf{u}(x, y, 0) = \mathbf{u}_0(x, y)\tag{2.7}$$

and the following boundary conditions:

$$\boldsymbol{\nu} \cdot \mathbf{u} = 0, \quad \boldsymbol{\nu} \cdot \nabla (\mathbf{u} \cdot \boldsymbol{\tau}) = 0, \quad (\alpha n \nabla c - \nabla n) \cdot \boldsymbol{\nu} = 0, \quad c = 1, \quad \forall (x, y) \in \Gamma,\tag{2.8}$$

$$\mathbf{u} = \mathbf{0}, \quad \nabla n \cdot \boldsymbol{\nu} = 0, \quad \nabla c \cdot \boldsymbol{\nu} = 0, \quad \forall (x, y) \in \partial\Omega_{\text{bot}}.\tag{2.9}$$

3. Diffuse-Domain Reformulation

3.1. Chemotaxis-fluid diffuse-domain model

In order to numerically solve the coupled chemotaxis-fluid system (2.3)–(2.9) in the drop domain, we propose a diffuse domain approximation of the chemotaxis-fluid model in a larger rectangular domain $\tilde{\Omega}$ outlined in Fig. 2. The cf-DD model reads as

$$\phi n_t + \nabla \cdot (\phi \mathbf{u}n) + \alpha \nabla \cdot [r(c)\phi n \nabla c] = \nabla \cdot (\phi \nabla n) + \mathcal{B}_n,\tag{3.1}$$

$$\phi c_t + \phi \mathbf{u} \cdot \nabla c = \delta \nabla \cdot (\phi \nabla c) - \beta r(c)\phi n + \mathcal{B}_c,\tag{3.2}$$

$$\phi \mathbf{u}_t + \phi \mathbf{u} \cdot \nabla \mathbf{u} + \text{Sc} \phi \nabla p = \text{Sc} \nabla \cdot (\phi \nabla \mathbf{u}) - \text{Sc} \gamma \phi n \mathbf{z} + \mathcal{B}_u,\tag{3.3}$$

$$\nabla \cdot (\phi \mathbf{u}) = 0,\tag{3.4}$$

with the boundary conditions

$$\boldsymbol{\nu} \cdot \nabla \mathbf{u} = \mathbf{0}, \quad (\alpha n \nabla c - \nabla n) \cdot \boldsymbol{\nu} = 0, \quad c = 1, \quad \forall (x, y) \in \tilde{\Gamma}, \quad (3.5)$$

$$\mathbf{u} = \mathbf{0}, \quad \nabla n \cdot \boldsymbol{\nu} = 0, \quad \nabla c \cdot \boldsymbol{\nu} = 0, \quad \forall (x, y) \in \partial \tilde{\Omega}_{\text{bot}}. \quad (3.6)$$

We set the diffuse-domain function ϕ to be the following approximation of the characteristic function of the original domain Ω :

$$\phi(\mathbf{x}) = \frac{1}{2} \left[1 - \tanh \left(\frac{3d(\mathbf{x})}{\varepsilon} \right) \right], \quad (3.7)$$

where $d(\mathbf{x})$ is the signed distance function to Γ ($d < 0$ inside Ω) and ε is the thickness of the diffuse domain boundary as shown in Fig. 2. Note that the function ϕ is independent of time since the domain Ω is fixed. Finally, the terms

$$\mathcal{B}_n = 0, \quad \mathcal{B}_c = -\frac{1-\phi}{\varepsilon^3}(c-1), \quad \mathcal{B}_u = \mathbf{0} \quad (3.8)$$

are added to enforce the original boundary conditions (2.8) on Γ , and these terms have been selected following the ideas introduced in Ref. 35. In Sec. 3.2, we will show that the cf-DD system (3.1)–(3.8) asymptotically converges to the original chemotaxis-fluid system (2.3)–(2.6) with the boundary conditions (2.8) and (2.9) as $\varepsilon \rightarrow 0$.




Fig. 2. (Color online) Schematic representation of the diffuse-domain method. The original domain Ω is embedded in a larger, rectangular domain $\tilde{\Omega}$, where a diffuse-domain function ϕ approximates the characteristic function of Ω . The boundary conditions on $\partial \tilde{\Omega}_{\text{bot}}$ are the same as those prescribed for the original system on $\partial \Omega_{\text{bot}}$, while the boundary conditions on $\tilde{\Gamma}$ are chosen to be consistent with those on Γ .

3.2. Asymptotic analysis

We now use the method of matched asymptotic expansions (see, e.g. Refs. 4, 10, 22, 26 and 39) to analyze the cf-DD system (3.1)–(3.8). In particular, we expand n , c , u , v , and p with respect to the small parameter ε (representing the interface thickness according to (3.7)) in regions close to the interface (inner region) and far from the interface (outer region), which are defined as follows: inner region $= \{\mathbf{x} : |d(\mathbf{x})| < \varepsilon^{s_1}\}$, outer region $= \{\mathbf{x} : |d(\mathbf{x})| > \varepsilon^{s_2}\}$, where $0 < s_1 < s_2$. When ε is small, the inner and outer regions overlap, and the two expansions are to be matched in the overlapping region $= \{\mathbf{x} : \varepsilon^{s_2} < |d(\mathbf{x})| < \varepsilon^{s_1}\}$.

For the purpose of asymptotic analysis, we consider a smoothed $r(c)$, while in the numerical experiments reported in Sec. 5 the original formula (1.2) has been utilized.

3.2.1. Outer expansions

We introduce the vector $\mathbf{w} := (n, c, u, v, p)^\top$ and expand it in ε in the outer region on each side of the interface Γ . We denote these formal outer expansions by $\mathbf{w}^+(\mathbf{x})$ for $\mathbf{x} \in \Omega$ at which $\phi(\mathbf{x}) \approx 1$ (inside Ω) and $\mathbf{w}^-(\mathbf{x})$ for $\mathbf{x} \notin \Omega$ at which $\phi(\mathbf{x}) \approx 0$ (outside Ω):

$$\mathbf{w}^\pm = \mathbf{w}_0^\pm + \varepsilon \mathbf{w}_1^\pm + \varepsilon^2 \mathbf{w}_2^\pm + \dots$$

We first substitute \mathbf{w}^+ into (3.1)–(3.4). Taking into account that $\phi \approx 1$ inside Ω , we combine the leading $\mathcal{O}(1)$ terms in the resulting expansions and obtain

$$\begin{aligned} (n_0^+)_t + \nabla \cdot (\mathbf{u}_0^+ n_0^+) + \alpha \nabla \cdot [r(c_0^+) n_0^+ \nabla c_0^+] &= \Delta n_0^+, \\ (c_0^+)_t + \mathbf{u}_0^+ \cdot \nabla c_0^+ &= \delta \Delta c_0^+ - \beta r(c_0^+) n_0^+, \\ (\mathbf{u}_0^+)_t + \mathbf{u}_0^+ \cdot \nabla \mathbf{u}_0^+ + \text{Sc} \nabla p_0^+ &= \text{Sc} \Delta \mathbf{u}_0^+ - \text{Sc} \gamma n_0^+ \mathbf{z}, \\ \nabla \cdot \mathbf{u}_0^+ &= 0, \end{aligned} \tag{3.9}$$

so that n_0^+ , c_0^+ , \mathbf{u}_0^+ , and p_0^+ satisfy the chemotaxis-fluid system (2.3)–(2.6). Moreover, substituting \mathbf{w}^+ into the boundary condition (3.6), we can easily see that n_0^+ , c_0^+ , and \mathbf{u}_0^+ satisfy the boundary condition (2.9). We then perform similar analysis for \mathbf{w}^- , which results in $c_0^- \equiv 1$ since $\phi \approx 0$ outside Ω .

3.2.2. Inner expansions

We now consider the expansions of \mathbf{w} in the inner region. To this end, we first use the divergence-free condition (3.4) and rewrite the system (3.1)–(3.4), (3.8) in the following equivalent form:

$$\phi n_t + \phi \mathbf{u} \cdot \nabla n + \alpha \nabla \cdot [r(c) \phi n \nabla c] = \nabla \cdot (\phi \nabla n), \tag{3.10}$$

$$\phi c_t + \phi \mathbf{u} \cdot \nabla c = \delta \nabla \cdot (\phi \nabla c) - \beta r(c) \phi n - (1 - \phi) \varepsilon^{-3} (c - 1), \tag{3.11}$$

$$\phi \mathbf{u}_t + \phi \mathbf{u} \cdot \nabla \mathbf{u} + \text{Sc} \phi \nabla p = \text{Sc} \nabla \cdot (\phi \nabla \mathbf{u}) - \text{Sc} \gamma \phi n \mathbf{z}, \quad (3.12)$$

$$\nabla \cdot (\phi \mathbf{u}) = 0. \quad (3.13)$$

Next, we introduce a rescaled variable $\xi := d(\mathbf{x})/\varepsilon$ and a local coordinate system near the interface Γ : $\mathbf{x}(s, \xi; \varepsilon) = \mathbf{X}(s) + \varepsilon \xi \boldsymbol{\nu}(s)$, where $\mathbf{X}(s)$ is a parametrization of Γ , and s is the arc length parameter. We use the notation $\omega(\mathbf{x}) = \widehat{\omega}(s, \xi)$ for any function ω , and notice that the following identities hold:

$$\nabla \omega = (1 + \varepsilon \xi \kappa)^{-1} \nabla_\Gamma \widehat{\omega} + \varepsilon^{-1} \boldsymbol{\nu} \widehat{\omega}_\xi, \quad (3.14)$$

$$\Delta \omega = (1 + \varepsilon \xi \kappa)^{-1} \nabla_\Gamma \cdot ((1 + \varepsilon \xi \kappa)^{-1} \nabla_\Gamma \widehat{\omega}) + \varepsilon^{-1} \kappa (1 + \varepsilon \xi \kappa)^{-1} \widehat{\omega}_\xi + \varepsilon^{-2} \widehat{\omega}_{\xi\xi}, \quad (3.15)$$

where, the ∇_Γ and $\nabla_\Gamma \cdot$ stand for the curve gradient and curve divergence operators, respectively. In (3.15), we have used the facts that $\nabla_\Gamma \cdot \boldsymbol{\nu} = \kappa$, where κ is the mean curvature of the interface, and $\nabla d = \boldsymbol{\nu}$.

We then substitute a formal expansion, which is valid in the inner region,

$$\widehat{\omega} = \widehat{\omega}_0 + \varepsilon \widehat{\omega}_1 + \varepsilon^2 \widehat{\omega}_2 + \dots \quad (3.16)$$

Into the system (3.10)–(3.13), use relations (3.14)–(3.15), and collect the like powers of ε . At the leading order term, $\mathcal{O}(\varepsilon^{-3})$, we obtain

$$(1 - \phi)(\widehat{c}_0 - 1) = 0. \quad (3.17)$$

Next, equating the $\mathcal{O}(\varepsilon^{-2})$ terms in Eqs. (3.10)–(3.12) results in

$$[\phi \alpha r(\widehat{c}_0) \widehat{n}_0(\widehat{c}_0)_\xi - \phi(\widehat{n}_0)_\xi]_\xi = 0, \quad (3.18)$$

$$\delta[\phi(\widehat{c}_0)_\xi]_\xi - (1 - \phi)\widehat{c}_1 = 0, \quad (3.19)$$

$$[\phi(\widehat{\mathbf{u}}_0)_\xi]_\xi = 0. \quad (3.20)$$

Finally, balancing the $\mathcal{O}(\varepsilon^{-1})$ terms in Eqs. (3.10)–(3.13) leads, after some simplifications, to

$$\begin{aligned} & \phi \widehat{\mathbf{u}}_0 \cdot \boldsymbol{\nu}(\widehat{n}_0)_\xi + [\phi \alpha r(\widehat{c}_0) \{ \widehat{n}_0(\widehat{c}_1)_\xi + \widehat{n}_1(\widehat{c}_0)_\xi \} + \phi \alpha r'(\widehat{c}_0) \widehat{c}_1 \widehat{n}_0(\widehat{c}_0)_\xi - \phi(\widehat{n}_1)_\xi]_\xi \\ & + \kappa \phi [\alpha r(\widehat{c}_0) \widehat{n}_0(\widehat{c}_0)_\xi - (\widehat{n}_0)_\xi] = 0, \end{aligned} \quad (3.21)$$

$$\phi \widehat{\mathbf{u}}_0 \cdot \boldsymbol{\nu}(\widehat{c}_0)_\xi + \delta[\phi(\widehat{c}_1)_\xi]_\xi + \delta \kappa \phi(\widehat{c}_0)_\xi - (1 - \phi)\widehat{c}_2 = 0, \quad (3.22)$$

$$\begin{aligned} & \phi \widehat{\mathbf{u}}_0 \cdot \boldsymbol{\nu}(\widehat{\mathbf{u}}_0)_\xi + \text{Sc} \{ \phi(\widehat{p}_0)_\xi \boldsymbol{\nu} - [\phi(\widehat{\mathbf{u}}_1)_\xi]_\xi - (\phi \boldsymbol{\nu} \cdot \nabla_\Gamma \widehat{\mathbf{u}}_0)_\xi \\ & - \nabla_\Gamma \cdot (\phi \boldsymbol{\nu})(\widehat{\mathbf{u}}_0)_\xi \} = \mathbf{0}, \end{aligned} \quad (3.23)$$

$$\boldsymbol{\nu} \cdot (\phi \widehat{\mathbf{u}}_0)_\xi = 0. \quad (3.24)$$

3.2.3. Matching conditions in the overlapping region

In what follows, we derive the boundary conditions on Γ by matching the outer and inner expansions in the overlapping region. To this end, the following matching conditions at $\mathbf{x} = \mathbf{X}(s) \in \Gamma$ need to be satisfied (see Refs. 1 and 22):

$$\lim_{\xi \rightarrow \mp\infty} \widehat{\mathbf{w}}_0(s, \xi) = \mathbf{w}_0^\pm(\mathbf{x}\pm), \quad (3.25)$$

$$\lim_{\xi \rightarrow \mp\infty} (\widehat{\mathbf{w}}_1(s, \xi))_\xi = \boldsymbol{\nu} \cdot \nabla \mathbf{w}_0^\pm(\mathbf{x}\pm), \quad (3.26)$$

where

$$\mathbf{w}_0^\pm(\mathbf{x}\pm) := \lim_{h \rightarrow 0^-} \mathbf{w}_0^\pm(\mathbf{x} \pm h\boldsymbol{\nu}).$$

First, we note that (3.17) implies $\widehat{c}_0(s, \xi) \equiv 1$, which together with the matching condition (3.25) imply

$$c_0^+(\mathbf{x}+) = \widehat{c}_0 \equiv 1 \quad \text{at } \mathbf{x} \in \Gamma. \quad (3.27)$$

We then use (3.27) to rewrite (3.18) as $[\phi(\widehat{n}_0)_\xi]_\xi = 0$, which implies $\phi(\widehat{n}_0)_\xi = C(s)$, and since $\lim_{\xi \rightarrow \infty} \phi = 0$, we conclude that $C(s) \equiv 0$, and hence $(\widehat{n}_0)_\xi = 0$. This together with (3.27) allows us to rewrite (3.21) as

$$[\phi(\alpha\widehat{n}_0(\widehat{c}_1)_\xi - (\widehat{n}_1)_\xi)]_\xi = 0,$$

and therefore, $\alpha\widehat{n}_0(\widehat{c}_1)_\xi - (\widehat{n}_1)_\xi = 0$, which, using the matching conditions (3.25) and (3.26), reduces to the following condition on n_0^+ and c_0^+ :

$$(\alpha n_0^+ \nabla c_0^+ - \nabla n_0^+) \cdot \boldsymbol{\nu} = 0. \quad (3.28)$$

Similarly, we use (3.20) to obtain $(\widehat{\mathbf{u}}_0)_\xi = \mathbf{0}$, which allows us to rewrite (3.23) as

$$\phi(\widehat{p}_0)_\xi \boldsymbol{\nu} - [\phi(\widehat{\mathbf{u}}_1)_\xi]_\xi = \mathbf{0}.$$

After applying the orthogonal projection operator $P_\Gamma := I - \boldsymbol{\nu} \otimes \boldsymbol{\nu} : \widetilde{\Omega} \rightarrow \Gamma$, where I is the identity matrix, the last equation further reduces to $[\phi(\widehat{\mathbf{u}}_1 \cdot \boldsymbol{\tau})_\xi]_\xi = 0$. We then proceed with the arguments similar to those used to derive (3.28) and conclude with

$$\boldsymbol{\nu} \cdot \nabla (\mathbf{u}_0^+ \cdot \boldsymbol{\tau}) = 0. \quad (3.29)$$

Also note that Eq. (3.24) and the matching condition (3.25) lead to

$$\boldsymbol{\nu} \cdot \mathbf{u}_0^+ = 0. \quad (3.30)$$

Finally, (3.27)–(3.30) together with (3.9) imply that the cf-DD system (3.1)–(3.8) asymptotically converges to the chemotaxis-fluid system (2.3)–(2.6) with the boundary conditions (2.8)–(2.9) as $\varepsilon \rightarrow 0$.

4. Hybrid Finite-Volume Finite-Difference Method

Recall that the advantage of the cf-DD system (3.1)–(3.8) is that it is posed on a simple, rectangular domain and thus it can be numerically solved in a much easier way compared to the original chemotaxis-fluid system. In this section, we provide a detailed description of the hybrid finite-volume finite-difference numerical method used to solve the studied cf-DD system.

The cell density equation (3.1) will be solved using a semi-discrete second-order finite-volume upwind scheme combined with a second-order strong stability-preserving multistep ODE solver for the temporal discretization. The oxygen concentration equation (3.2) and the Navier–Stokes fluid equations will be discretized using a semi-discrete finite-difference central scheme combined with a second-order BDF-like method for time integration and a second-order projection method needed to enforce the divergence-free property of the fluid velocity.

4.1. Finite-volume upwind scheme for the cell density equation

We first define $m := \phi n$, substitute (3.8) into the cell density equation (3.1) and rewrite it in an equivalent coordinate form:

$$m_t + [(u + \alpha r(c)c_x)m]_x + [(v + \alpha r(c)c_y)m]_y = \left[\phi \left(\frac{m}{\phi} \right)_x \right]_x + \left[\phi \left(\frac{m}{\phi} \right)_y \right]_y. \quad (4.1)$$

We then discretize equation (4.1) in space using the semi-discrete second-order finite-volume upwind scheme from Ref. 15.

To this end, we divide the computational domain $\tilde{\Omega}$ into the Cartesian cells $I_{j,k} := [x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}] \times [y_{k-\frac{1}{2}}, y_{k+\frac{1}{2}}]$, centered at $(x_j, y_k) = (\frac{x_{j-\frac{1}{2}} + x_{j+\frac{1}{2}}}{2}, \frac{y_{k-\frac{1}{2}} + y_{k+\frac{1}{2}}}{2})$ with $j = 1, \dots, N$ and $k = 1, \dots, M$. In principle, this mesh can consist of the rectangles of different size, but for simplicity of presentation, we use a uniform mesh with $x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}} \equiv \Delta x$ and $y_{k+\frac{1}{2}} - y_{k-\frac{1}{2}} \equiv \Delta y$, where Δx and Δy are (small) constants. We then denote the cell averages of m by

$$\bar{m}_{j,k}(t) \approx \frac{1}{\Delta x \Delta y} \iint_{I_{j,k}} m(x, y, t) dx dy,$$

and integrate equation (4.1) over cell $I_{j,k}$ to obtain

$$\begin{aligned} & \Delta x \Delta y \frac{d}{dt} \bar{m}_{j,k}(t) + \int_{y_{k-\frac{1}{2}}}^{y_{k+\frac{1}{2}}} (u + \alpha r(c)c_x) m \Big|_{x_{j-\frac{1}{2}}}^{x_{j+\frac{1}{2}}} dy \\ & + \int_{x_{j-\frac{1}{2}}}^{x_{j+\frac{1}{2}}} (v + \alpha r(c)c_y) m \Big|_{y_{k-\frac{1}{2}}}^{y_{k+\frac{1}{2}}} dx \\ & = \int_{y_{k-\frac{1}{2}}}^{y_{k+\frac{1}{2}}} \phi \left(\frac{m}{\phi} \right)_x \Big|_{x_{j-\frac{1}{2}}}^{x_{j+\frac{1}{2}}} dy + \int_{x_{j-\frac{1}{2}}}^{x_{j+\frac{1}{2}}} \phi \left(\frac{m}{\phi} \right)_y \Big|_{y_{k-\frac{1}{2}}}^{y_{k+\frac{1}{2}}} dx. \end{aligned}$$

Applying the midpoint rule to the above integrals and dividing by $\Delta x \Delta y$ results in

$$\begin{aligned} \frac{d}{dt} \bar{m}_{j,k} = & - \frac{(u + \alpha r(c)c_x)m|_{(x_{j+\frac{1}{2}}, y_k)} - (u + \alpha r(c)c_x)m|_{(x_{j-\frac{1}{2}}, y_k)}}{\Delta x} \\ & - \frac{(v + \alpha r(c)c_y)m|_{(x_j, y_{k+\frac{1}{2}})} - (v + \alpha r(c)c_y)m|_{(x_j, y_{k-\frac{1}{2}})}}{\Delta y} \\ & + \frac{\phi\left(\frac{m}{\phi}\right)_x|_{(x_{j+\frac{1}{2}}, y_k)} - \phi\left(\frac{m}{\phi}\right)_x|_{(x_{j-\frac{1}{2}}, y_k)}}{\Delta x} \\ & + \frac{\phi\left(\frac{m}{\phi}\right)_y|_{(x_j, y_{k+\frac{1}{2}})} - \phi\left(\frac{m}{\phi}\right)_y|_{(x_j, y_{k-\frac{1}{2}})}}{\Delta y}. \end{aligned} \quad (4.2)$$

We note that $\bar{m}_{j,k}$ as well as many other indexed quantities in (4.2) and below depend on time t , but from now on we omit this dependence for the sake of brevity.

The construction of the scheme will be completed once the fluxes at the cell interfaces in (4.2) are approximated numerically. The semi-discrete finite-volume upwind scheme can then be written as the following system of ODEs:

$$\begin{aligned} \frac{d}{dt} \bar{m}_{j,k} = & - \frac{F_{j+\frac{1}{2},k}^x - F_{j-\frac{1}{2},k}^x}{\Delta x} - \frac{F_{j,k+\frac{1}{2}}^y - F_{j,k-\frac{1}{2}}^y}{\Delta y} \\ & + \frac{G_{j+\frac{1}{2},k}^x - G_{j-\frac{1}{2},k}^x}{\Delta x} + \frac{G_{j,k+\frac{1}{2}}^y - G_{j,k-\frac{1}{2}}^y}{\Delta y}, \end{aligned} \quad (4.3)$$

where

$$F_{j\pm\frac{1}{2},k}^x \approx (u + \alpha r(c)c_x)m|_{(x_{j\pm\frac{1}{2}}, y_k)} \quad \text{and} \quad F_{j,k\pm\frac{1}{2}}^y \approx (v + \alpha r(c)c_y)m|_{(x_j, y_{k\pm\frac{1}{2}})} \quad (4.4)$$

are numerical convection–chemotaxis fluxes, and

$$G_{j\pm\frac{1}{2},k}^x \approx \phi\left(\frac{m}{\phi}\right)_x|_{(x_{j\pm\frac{1}{2}}, y_k)} \quad \text{and} \quad G_{j,k\pm\frac{1}{2}}^y \approx \phi\left(\frac{m}{\phi}\right)_y|_{(x_j, y_{k\pm\frac{1}{2}})} \quad (4.5)$$

are centered numerical diffusion fluxes.

In order to ensure stability of the scheme (4.3)–(4.5), we use an upwind approximation of the convection–chemotaxis fluxes, which can be written in the following form:

$$\begin{aligned} F_{j+\frac{1}{2},k}^x &= \begin{cases} a_{j+\frac{1}{2},k} m_{j,k}^E & \text{if } a_{j+\frac{1}{2},k} \geq 0, \\ a_{j+\frac{1}{2},k} m_{j+1,k}^W & \text{if } a_{j+\frac{1}{2},k} < 0, \end{cases} \\ F_{j,k+\frac{1}{2}}^y &= \begin{cases} b_{j,k+\frac{1}{2}} m_{j,k}^N & \text{if } b_{j,k+\frac{1}{2}} \geq 0, \\ b_{j,k+\frac{1}{2}} m_{j,k+1}^S & \text{if } b_{j,k+\frac{1}{2}} < 0. \end{cases} \end{aligned} \quad (4.6)$$

Here, $m_{j,k}^{E,W,N,S}$ are the point values of the piecewise linear reconstruction consisting of the following linear pieces on every interval $I_{j,k}$:

$$\tilde{m}_{j,k}(x, y) = \bar{m}_{j,k} + (m_x)_{j,k}(x - x_j) + (m_y)_{j,k}(y - y_k), \quad (x, y) \in I_{j,k}, \quad (4.7)$$

at the points $(x_{j+\frac{1}{2}}, y_k)$, $(x_{j-\frac{1}{2}}, y_k)$, $(x_j, y_{k+\frac{1}{2}})$, and $(x_j, y_{k-\frac{1}{2}})$, respectively. Namely, we have

$$\begin{aligned} m_{j,k}^E &= \tilde{m}_{j,k}(x_{j+\frac{1}{2}}, y_k) = \bar{m}_{j,k} + \frac{\Delta x}{2}(m_x)_{j,k}, \\ m_{j,k}^W &= \tilde{m}_{j,k}(x_{j-\frac{1}{2}}, y_k) = \bar{m}_{j,k} - \frac{\Delta x}{2}(m_x)_{j,k}, \\ m_{j,k}^N &= \tilde{m}_{j,k}(x_j, y_{k+\frac{1}{2}}) = \bar{m}_{j,k} + \frac{\Delta y}{2}(m_y)_{j,k}, \\ m_{j,k}^S &= \tilde{m}_{j,k}(x_j, y_{k-\frac{1}{2}}) = \bar{m}_{j,k} - \frac{\Delta y}{2}(m_y)_{j,k}. \end{aligned} \quad (4.8)$$

The second order of accuracy will be guaranteed provided the numerical derivatives $(m_x)_{j,k}$ and $(m_y)_{j,k}$ are to be (at least) first-order approximations of the corresponding exact derivatives $m_x(x_j, y_k, t)$ and $m_y(x_j, y_k, t)$. In our numerical experiments, we have used the central-difference approximations

$$(m_x)_{j,k} = \frac{\bar{m}_{j+1,k} - \bar{m}_{j-1,k}}{2\Delta x} \quad \text{and} \quad (m_y)_{j,k} = \frac{\bar{m}_{j,k+1} - \bar{m}_{j,k-1}}{2\Delta y}, \quad (4.9)$$

throughout the computational domain except for the cells, where the linear approach (4.9) leads to the appearance of negative reconstructed values of m in (4.8). In the cells, where either $m_{j,k}^E$ or $m_{j,k}^W$ is negative, we replace (4.9) with a nonlinear minmod2 reconstruction (see, e.g. Refs. 37, 41 and 45):

$$(m_x)_{j,k} = \text{minmod}\left(2 \frac{\bar{m}_{j,k} - \bar{m}_{j-1,k}}{\Delta x}, \frac{\bar{m}_{j+1,k} - \bar{m}_{j-1,k}}{2\Delta x}, 2 \frac{\bar{m}_{j+1,k} - \bar{m}_{j,k}}{\Delta x}\right), \quad (4.10)$$

which guarantees that no negative values of m emerge in (4.8). We then recalculate the reconstructed values $m_{j,k}^E$ and $m_{j,k}^W$. Similarly, if either $m_{j,k}^N$ or $m_{j,k}^S$ is negative, we set

$$(m_y)_{j,k} = \text{minmod}\left(2 \frac{\bar{m}_{j,k} - \bar{m}_{j,k-1}}{\Delta y}, \frac{\bar{m}_{j,k+1} - \bar{m}_{j,k-1}}{2\Delta y}, 2 \frac{\bar{m}_{j,k+1} - \bar{m}_{j,k}}{\Delta y}\right), \quad (4.11)$$

and recalculate the reconstructed values $m_{j,k}^N$ and $m_{j,k}^S$. The minmod function used in (4.10) and (4.11) is defined as

$$\text{minmod}(z_1, z_2, \dots) := \begin{cases} \min_j \{z_j\} & \text{if } z_j > 0 \ \forall j, \\ \max_j \{z_j\} & \text{if } z_j < 0 \ \forall j, \\ 0 & \text{otherwise.} \end{cases} \quad (4.12)$$

The description of the numerical convection–chemotaxis fluxes in (4.6) will be completed once the local speeds in the x - and y -directions, $a_{j+\frac{1}{2},k}$ and $b_{j,k+\frac{1}{2}}$, are specified. Since all of the solution components are expected to be smooth, the local speeds can be approximated using the centered differences and averages as

$$\begin{aligned} a_{j+\frac{1}{2},k} &= u_{j+\frac{1}{2},k} + \alpha r(c_{j+\frac{1}{2},k})(c_x)_{j+\frac{1}{2},k} \quad \text{and} \\ b_{j,k+\frac{1}{2}} &= v_{j,k+\frac{1}{2}} + \alpha r(c_{j,k+\frac{1}{2}})(c_y)_{j,k+\frac{1}{2}}, \end{aligned}$$

where

$$\begin{aligned} (c_x)_{j+\frac{1}{2},k} &= \frac{c_{j+1,k} - c_{j,k}}{\Delta x}, \quad u_{j+\frac{1}{2},k} = \frac{1}{2}(u_{j,k}^E + u_{j+1,k}^W), \quad c_{j+\frac{1}{2},k} = \frac{1}{2}(c_{j,k}^E + c_{j+1,k}^W), \\ (c_y)_{j,k+\frac{1}{2}} &= \frac{c_{j,k+1} - c_{j,k}}{\Delta y}, \quad v_{j,k+\frac{1}{2}} = \frac{1}{2}(v_{j,k}^N + v_{j,k+1}^S), \quad c_{j,k+\frac{1}{2}} = \frac{1}{2}(c_{j,k}^N + c_{j,k+1}^S). \end{aligned}$$

Here, the point values $c_{j,k}^{E,W,N,S}$, $u_{j,k}^{E,W}$, and $v_{j,k}^{N,S}$ are obtained using the same piecewise linear reconstruction, which was used to compute the corresponding values of m in (4.8), but now applied to the point values $c_{j,k} \approx c(x_j, y_k, t)$, $u_{j,k} \approx u(x_j, y_k, t)$, and $v_{j,k} \approx v(x_j, y_k, t)$, respectively.

Finally, the centered numerical diffusion fluxes in (4.5) are approximated by

$$\begin{aligned} G_{j+\frac{1}{2},k}^x &= \frac{\phi_{j+\frac{1}{2},k}}{\Delta x} \left(\frac{\bar{m}_{j+1,k}}{\phi_{j+1,k}} - \frac{\bar{m}_{j,k}}{\phi_{j,k}} \right), \\ G_{j,k+\frac{1}{2}}^y &= \frac{\phi_{j,k+\frac{1}{2}}}{\Delta y} \left(\frac{\bar{m}_{j,k+1}}{\phi_{j,k+1}} - \frac{\bar{m}_{j,k}}{\phi_{j,k}} \right), \end{aligned} \quad (4.13)$$

where $\phi_{j,k} := \phi(x_j, y_k)$, $\phi_{j+\frac{1}{2},k} := \phi(x_{j+\frac{1}{2}}, y_k)$, and $\phi_{j,k+\frac{1}{2}} := \phi(x_j, y_{k+\frac{1}{2}})$.

Time discretization. The semi-discretization (4.3) results in the system of time-dependent ODEs, which we integrate using the second-order strong stability-preserving (SSP) three-step method Ref. 19. This results in

$$\begin{aligned} \bar{m}_{j,k}^{\ell+1} &= \frac{3}{4} \bar{m}_{j,k}^\ell - \frac{3}{2} \lambda \left(F_{j+\frac{1}{2},k}^{x,\ell} - F_{j-\frac{1}{2},k}^{x,\ell} \right) - \frac{3}{2} \mu \left(F_{j,k+\frac{1}{2}}^{y,\ell} - F_{j,k-\frac{1}{2}}^{y,\ell} \right) \\ &\quad + \frac{3}{2} \lambda \left(G_{j+\frac{1}{2},k}^{x,\ell} - G_{j-\frac{1}{2},k}^{x,\ell} \right) + \frac{3}{2} \mu \left(G_{j,k+\frac{1}{2}}^{y,\ell} - G_{j,k-\frac{1}{2}}^{y,\ell} \right) + \frac{1}{4} \bar{m}_{j,k}^{\ell-2}, \end{aligned} \quad (4.14)$$

where Δt is the time step, $\lambda := \Delta t / \Delta x$, $\mu := \Delta t / \Delta y$, $t^\ell := \ell \Delta t$, $\bar{m}_{j,k}^\ell := \bar{m}_{j,k}(t^\ell)$, $F_{j+\frac{1}{2},k}^{x,\ell} := F_{j+\frac{1}{2},k}^x(t^\ell)$, $F_{j+\frac{1}{2},k}^{y,\ell} := F_{j+\frac{1}{2},k}^y(t^\ell)$, $G_{j,k+\frac{1}{2}}^{x,\ell} := G_{j,k+\frac{1}{2}}^x(t^\ell)$, and $G_{j,k+\frac{1}{2}}^{y,\ell} := G_{j,k+\frac{1}{2}}^y(t^\ell)$.

The resulting fully discrete scheme (4.14) is positivity preserving in the sense that $\bar{m}_{j,k}^{\ell+1} \geq 0$ for all j, k provided $\bar{m}_{j,k}^\ell \geq 0$ and $\bar{m}_{j,k}^{\ell-2} \geq 0$ for all j, k and Δt is sufficiently small. In order to prove this, we first note that the convection–chemotaxis numerical fluxes (4.6) can be rewritten as

$$F_{j+\frac{1}{2},k}^{x,\ell} = a_{j+\frac{1}{2},k} \left(\frac{1 + \text{sign}(a_{j+\frac{1}{2},k})}{2} m_{j,k}^E + \frac{1 - \text{sign}(a_{j+\frac{1}{2},k})}{2} m_{j+1,k}^W \right) \quad (4.15)$$

and

$$F_{j,k+\frac{1}{2}}^{y,\ell} = b_{j,k+\frac{1}{2}} \left(\frac{1 + \text{sign}(b_{j,k+\frac{1}{2}})}{2} m_{j,k}^N + \frac{1 - \text{sign}(b_{j,k+\frac{1}{2}})}{2} m_{j,k+1}^S \right), \quad (4.16)$$

and by the conservation property of the piecewise-linear reconstruction (4.7) the identity

$$\bar{m}_{j,k}^\ell = \frac{1}{8} (m_{j,k}^E + m_{j,k}^W + m_{j,k}^N + m_{j,k}^S) + \frac{1}{2} \bar{m}_{j,k}^\ell \quad (4.17)$$

holds. Note that the quantities $a_{j+\frac{1}{2},k}$, $b_{j,k+\frac{1}{2}}$, and $m_{j,k}^{E,W,N,S}$ in (4.15)–(4.17) are evaluated at time level $t = t^\ell$. We then substitute (4.13) and (4.15)–(4.17) into (4.14) to obtain

$$\begin{aligned} \bar{m}_{j,k}^{\ell+1} = & \frac{3}{4} \left\{ \left[\frac{1}{8} - \lambda |a_{j+\frac{1}{2},k}| (1 + \text{sign}(a_{j+\frac{1}{2},k})) \right] m_{j,k}^E \right. \\ & + \lambda |a_{j+\frac{1}{2},k}| (1 - \text{sign}(a_{j+\frac{1}{2},k})) m_{j+1,k}^W \\ & + \left[\frac{1}{8} - \lambda |a_{j-\frac{1}{2},k}| (1 - \text{sign}(a_{j-\frac{1}{2},k})) \right] m_{j,k}^W \\ & + \lambda |a_{j-\frac{1}{2},k}| (1 + \text{sign}(a_{j-\frac{1}{2},k})) m_{j-1,k}^E \\ & + \left[\frac{1}{8} - \mu |b_{j,k+\frac{1}{2}}| (1 + \text{sign}(b_{j,k+\frac{1}{2}})) \right] m_{j,k}^N \\ & + \mu |b_{j,k+\frac{1}{2}}| (1 - \text{sign}(b_{j,k+\frac{1}{2}})) m_{j,k+1}^S \\ & + \left[\frac{1}{8} - \mu |b_{j,k-\frac{1}{2}}| (1 - \text{sign}(b_{j,k-\frac{1}{2}})) \right] m_{j,k}^S \\ & \left. + \mu |b_{j,k-\frac{1}{2}}| (1 + \text{sign}(b_{j,k-\frac{1}{2}})) m_{j,k-1}^N \right\} \\ & + \frac{3}{2} \left\{ \left(\frac{1}{4} - \Delta t \left[\frac{\phi_{j+\frac{1}{2},k} + \phi_{j-\frac{1}{2},k}}{\phi_{j,k}(\Delta x)^2} + \frac{\phi_{j,k+\frac{1}{2}} + \phi_{j,k-\frac{1}{2}}}{\phi_{j,k}(\Delta y)^2} \right] \right) \bar{m}_{j,k}^\ell \right. \\ & + \frac{\Delta t}{(\Delta x)^2} \left(\frac{\phi_{j+\frac{1}{2},k}}{\phi_{j+1,k}} \bar{m}_{j+1,k}^\ell + \frac{\phi_{j-\frac{1}{2},k}}{\phi_{j-1,k}} \bar{m}_{j-1,k}^\ell \right) \\ & \left. + \frac{\Delta t}{(\Delta y)^2} \left(\frac{\phi_{j,k+\frac{1}{2}}}{\phi_{j,k+1}} \bar{m}_{j,k+1}^\ell + \frac{\phi_{j,k-\frac{1}{2}}}{\phi_{j,k-1}} \bar{m}_{j,k-1}^\ell \right) \right\} + \frac{1}{4} \bar{m}_{j,k}^{\ell-2}. \quad (4.18) \end{aligned}$$

As one can see from (4.18), the new values $\{\bar{m}_{j,k}^{\ell+1}\}$ are linear combinations of the non-negative cell averages $\{\bar{m}_{j,k}^\ell\}$ and $\{\bar{m}_{j,k}^{\ell-2}\}$, and the reconstructed point value $\{m_{j,k}^{E,W,N,S}\}$, which are also non-negative since they are computed using the

positivity preserving piecewise linear reconstruction (4.8)–(4.12). Thus, as long as the following CFL condition is satisfied:

$$\Delta t \leq \frac{1}{16} \min \left\{ \frac{\Delta x}{a_{\max}}, \frac{\Delta y}{b_{\max}}, \frac{4\phi_{j,k}(\Delta x)^2(\Delta y)^2}{(\phi_{j,k+\frac{1}{2}} + \phi_{j,k-\frac{1}{2}})(\Delta x)^2 + (\phi_{j+\frac{1}{2},k} + \phi_{j-\frac{1}{2},k})(\Delta y)^2} \right\}, \quad (4.19)$$

where

$$a_{\max} := \max_{j,k} \{ |a_{j+\frac{1}{2},k}| \}, \quad b_{\max} := \max_{j,k} \{ |b_{j,k+\frac{1}{2}}| \}, \quad (4.20)$$

the linear combination in (4.18) is a convex combination, which implies the non-negativity of $\bar{m}_{j,k}^{\ell+1}$ for all j, k .

Finally, since $m = \phi n$ and $\phi(\mathbf{x}) > 0$ for all \mathbf{x} , we conclude that $\bar{n}_{j,k}^{\ell+1} \geq 0$ for all j, k .

Remark 4.1. Note that the inequality (4.19) should be satisfied at every time level $t = t^\ell$, but since we use the three-step time discretization method, we have to choose a fixed Δt at time $t = 0$, when the data require to be used to evaluate the maxima in (4.20) are not available yet. We therefore replace a_{\max} and b_{\max} in (4.19) with their *a priori* upper bounds, which should be valid for all t and can be obtained for any problem at hand.

Remark 4.2. We note that we obtain $\bar{n}_{j,k}^1$ and $\bar{n}_{j,k}^2$ at the first two time steps using the first-order forward Euler time discretization.

4.2. Second-order projection finite-difference method for the Navier–Stokes and oxygen equations

Equipped with the obtained values $\bar{n}_{j,k}^{\ell+1}$, we now construct a second-order projection finite-difference method for Eqs. (3.2)–(3.4) by following the approach from Refs. 21 and 40 proposed in the context of the Cahn–Hilliard–Navier–Stokes system.

We begin with the second-order time discretization of (3.2)–(3.4), which is based on the projection method and the BDF method with Adams–Bashforth extrapolation. Assuming that $n^{\ell+1} \approx n(x, y, t^{\ell+1})$, $c^\ell \approx c(x, y, t^\ell)$, $\mathbf{u}^\ell \approx \mathbf{u}(x, y, t^\ell)$, and $p^\ell \approx p(x, y, t^\ell)$ are available, we obtain $c^{\ell+1}$, $\mathbf{u}^{\ell+1}$, and $p^{\ell+1}$ by solving the following equations:

$$\begin{aligned} \phi \frac{3\tilde{\mathbf{u}}^{\ell+1} - 4\mathbf{u}^\ell + \mathbf{u}^{\ell-1}}{2\Delta t} + \phi \mathbf{u}^* \cdot \nabla \mathbf{u}^* + \text{Sc} \phi \nabla p^\ell \\ = \text{Sc} \nabla \cdot (\phi \nabla \tilde{\mathbf{u}}^{\ell+1}) + \text{Sc} \gamma \phi n^{\ell+1} \mathbf{z}, \end{aligned} \quad (4.21)$$

$$\frac{3(\mathbf{u}^{\ell+1} - \tilde{\mathbf{u}}^{\ell+1})}{2\Delta t} + \text{Sc} \nabla \psi^{\ell+1} = \mathbf{0}, \quad (4.22)$$

$$\nabla \cdot (\phi \mathbf{u}^{\ell+1}) = 0, \quad (4.23)$$

$$p^{\ell+1} = \psi^{\ell+1} + p^\ell, \quad (4.24)$$

$$\begin{aligned} & \phi \frac{3c^{\ell+1} - 4c^\ell + c^{\ell-1}}{2\Delta t} + \phi \mathbf{u}^{\ell+1} \cdot \nabla c^* \\ &= \delta \nabla \cdot (\phi \nabla c^{\ell+1}) - \beta r(c^*) \phi n^{\ell+1} - \frac{1}{\varepsilon^3} (1 - \phi)(c^{\ell+1} - 1), \end{aligned} \quad (4.25)$$

where $\mathbf{u}^* := 2\mathbf{u}^\ell - \mathbf{u}^{\ell-1}$, $c^* = 2c^\ell - c^{\ell-1}$, and $\psi^{\ell+1}$ is an auxiliary variable.

The scheme (4.21)–(4.25) is implemented in the following way. First, we solve the elliptic equation (4.21) for $\tilde{\mathbf{u}}^{\ell+1}$ subject to the boundary conditions specified in (3.5)–(3.6) for \mathbf{u} . We multiply both sides of (4.22) by ϕ , take the divergence of the result, and use the divergence-free condition (4.23) to obtain the elliptic equation on $\psi^{\ell+1}$

$$\nabla \cdot (\phi \nabla \psi^{\ell+1}) = \frac{3}{2\text{Sc} \Delta t} \nabla \cdot (\phi \tilde{\mathbf{u}}^{\ell+1}), \quad (4.26)$$

which is solved subject to the homogeneous Neumann boundary condition $\nabla \psi^{\ell+1} \cdot \boldsymbol{\nu} = 0$ prescribed on $\partial\tilde{\Omega}$. Next, we substitute the computed $\psi^{\ell+1}$ into (4.22) and (4.24) to obtain $\mathbf{u}^{\ell+1}$ and $p^{\ell+1}$. Finally, we substitute $\mathbf{u}^{\ell+1}$ and $n^{\ell+1}$ into (4.25) and obtain the elliptic equation for $c^{\ell+1}$, which is solved subject to the boundary condition specified in (3.5) and (3.6).

Remark 4.3. It is shown in Ref. 20 that the scheme (4.21)–(4.24) without the source term $\text{Sc} \gamma \phi n^{\ell+1} \mathbf{z}$ is unconditionally stable.

Remark 4.4. We note that we obtain \mathbf{u}^1 , p^1 , and c^1 at the first time step using the following first-order time discretization, which is based on the projection method and the backward Euler method:

$$\phi \frac{\tilde{\mathbf{u}}^{\ell+1} - \mathbf{u}^\ell}{\Delta t} + \phi \mathbf{u}^\ell \cdot \nabla \mathbf{u}^\ell + \text{Sc} \phi \nabla p^\ell = \text{Sc} \nabla \cdot (\phi \nabla \tilde{\mathbf{u}}^{\ell+1}) + \text{Sc} \gamma \phi n^{\ell+1} \mathbf{z},$$

$$\frac{\mathbf{u}^{\ell+1} - \tilde{\mathbf{u}}^{\ell+1}}{\Delta t} + \text{Sc} \nabla (p^{\ell+1} - p^\ell) = \mathbf{0},$$

$$\nabla \cdot (\phi \mathbf{u}^{\ell+1}) = 0,$$

$$\phi \frac{c^{\ell+1} - c^\ell}{\Delta t} + \phi \mathbf{u}^{\ell+1} \cdot \nabla c^\ell$$

$$= \delta \nabla \cdot (\phi \nabla c^{\ell+1}) - \beta r(c^\ell) \phi n^{\ell+1} - \frac{1}{\varepsilon^3} (1 - \phi)(c^{\ell+1} - 1).$$

Spatial discretization. We now denote the point values of \mathbf{u} , ψ , p , and c at the cell centers (x_j, y_k) at time level $t = t^\ell$ by $\mathbf{u}_{j,k}^\ell$, $\psi_{j,k}^\ell$, $p_{j,k}^\ell$, and $c_{j,k}^\ell$, respectively, and apply the second-order central difference approximations to construct a fully discrete scheme.

First, we discretize equation (4.21) and use the cell averages $\bar{n}_{j,k}^{\ell+1}$ obtained in Sec. 4.1 to update $\tilde{u}_{j,k}^{\ell+1}$ and $\tilde{v}_{j,k}^{\ell+1}$ by solving the linear systems

$$\begin{aligned} \phi_{j,k} & \left[\frac{3\tilde{u}_{j,k}^{\ell+1} - 4u_{j,k}^{\ell} + u_{j,k}^{\ell-1}}{2\Delta t} + u_{j,k}^* \frac{u_{j+1,k}^* - u_{j-1,k}^*}{2\Delta x} \right. \\ & \quad \left. + v_{j,k}^* \frac{u_{j,k+1}^* - u_{j,k-1}^*}{2\Delta y} + \text{Sc} \frac{p_{j+1,k}^{\ell} - p_{j-1,k}^{\ell}}{2\Delta x} \right] \\ & = \text{Sc} \frac{\phi_{j+\frac{1}{2},k}(\tilde{u}_{j+1,k}^{\ell+1} - \tilde{u}_{j,k}^{\ell+1}) - \phi_{j-\frac{1}{2},k}(\tilde{u}_{j,k}^{\ell+1} - \tilde{u}_{j-1,k}^{\ell+1})}{(\Delta x)^2} \\ & \quad + \text{Sc} \frac{\phi_{j,k+\frac{1}{2}}(\tilde{u}_{j,k+1}^{\ell+1} - \tilde{u}_{j,k}^{\ell+1}) - \phi_{j,k-\frac{1}{2}}(\tilde{u}_{j,k}^{\ell+1} - \tilde{u}_{j,k-1}^{\ell+1})}{(\Delta y)^2}, \end{aligned}$$

and

$$\begin{aligned} \phi_{j,k} & \left[\frac{3\tilde{v}_{j,k}^{\ell+1} - 4v_{j,k}^{\ell} + v_{j,k}^{\ell-1}}{2\Delta t} + u_{j,k}^* \frac{v_{j+1,k}^* - v_{j-1,k}^*}{2\Delta x} + v_{j,k}^* \frac{v_{j,k+1}^* - v_{j,k-1}^*}{2\Delta y} \right. \\ & \quad \left. + \text{Sc} \frac{p_{j,k+1}^{\ell} - p_{j,k-1}^{\ell}}{2\Delta y} - \text{Sc} \gamma n_{j,k}^{\ell+1} \right] \\ & = \text{Sc} \frac{\phi_{j+\frac{1}{2},k}(\tilde{v}_{j+1,k}^{\ell+1} - \tilde{v}_{j,k}^{\ell+1}) - \phi_{j-\frac{1}{2},k}(\tilde{v}_{j,k}^{\ell+1} - \tilde{v}_{j-1,k}^{\ell+1})}{(\Delta x)^2} \\ & \quad + \text{Sc} \frac{\phi_{j,k+\frac{1}{2}}(\tilde{v}_{j,k+1}^{\ell+1} - \tilde{v}_{j,k}^{\ell+1}) - \phi_{j,k-\frac{1}{2}}(\tilde{v}_{j,k}^{\ell+1} - \tilde{v}_{j,k-1}^{\ell+1})}{(\Delta y)^2}, \end{aligned}$$

for $\{\tilde{u}_{j,k}^{\ell+1}\}$ and $\{\tilde{v}_{j,k}^{\ell+1}\}$, respectively. We then discretize equation (4.26) and obtain $\psi_{j,k}^{\ell+1}$ by solving the linear system

$$\begin{aligned} & \frac{\phi_{j+\frac{1}{2},k}(\psi_{j+1,k}^{\ell+1} - \psi_{j,k}^{\ell+1}) - \phi_{j-\frac{1}{2},k}(\psi_{j,k}^{\ell+1} - \psi_{j-1,k}^{\ell+1})}{(\Delta x)^2} \\ & + \frac{\phi_{j,k+\frac{1}{2}}(\psi_{j,k+1}^{\ell+1} - \psi_{j,k}^{\ell+1}) - \phi_{j,k-\frac{1}{2}}(\psi_{j,k}^{\ell+1} - \psi_{j,k-1}^{\ell+1})}{(\Delta y)^2} \\ & = \frac{3}{2\text{Sc} \Delta t} \left(\frac{\phi_{j+1,k}\tilde{u}_{j+1,k}^{\ell+1} - \phi_{j-1,k}\tilde{u}_{j-1,k}^{\ell+1}}{2\Delta x} + \frac{\phi_{j,k+1}\tilde{v}_{j,k+1}^{\ell+1} - \phi_{j,k-1}\tilde{v}_{j,k-1}^{\ell+1}}{2\Delta y} \right). \end{aligned}$$

Next, we find $u_{j,k}^{\ell+1}$, $v_{j,k}^{\ell+1}$, and $p_{j,k}^{\ell+1}$ by discretizing (4.22) and (4.24) as follows:

$$u_{j,k}^{\ell+1} = \tilde{u}_{j,k}^{\ell+1} - \frac{2}{3}\text{Sc} \Delta t \frac{\psi_{j+1,k}^{\ell+1} - \psi_{j-1,k}^{\ell+1}}{2\Delta x},$$

$$v_{j,k}^{\ell+1} = \tilde{v}_{j,k}^{\ell+1} - \frac{2}{3} \text{Sc} \Delta t \frac{\psi_{j,k+1}^{\ell+1} - \psi_{j,k-1}^{\ell+1}}{2\Delta y},$$

$$p_{j,k}^{\ell+1} = \psi_{j,k}^{\ell+1} + p_{j,k}^{\ell}.$$

Finally, we discretize (4.25) and update $c_{j,k}^{\ell+1}$ by solving the linear system

$$\begin{aligned} \phi_{j,k} & \left[\frac{3c_{j,k}^{\ell+1} - 4c_{j,k}^{\ell} + c_{j,k}^{\ell-1}}{2\Delta t} + u_{j,k}^{\ell+1} \frac{c_{j+1,k}^* - c_{j-1,k}^*}{2\Delta x} + v_{j,k}^{\ell+1} \frac{c_{j,k+1}^* - c_{j,k-1}^*}{2\Delta y} \right] \\ & = \delta \frac{\phi_{j+\frac{1}{2},k}(c_{j+1,k}^{\ell+1} - c_{j,k}^{\ell+1}) - \phi_{j-\frac{1}{2},k}(c_{j,k}^{\ell+1} - c_{j-1,k}^{\ell+1})}{(\Delta x)^2} \\ & \quad + \delta \frac{\phi_{j,k+\frac{1}{2}}(c_{j,k+1}^{\ell+1} - c_{j,k}^{\ell+1}) - \phi_{j,k-\frac{1}{2}}(c_{j,k}^{\ell+1} - c_{j,k-1}^{\ell+1})}{(\Delta y)^2} \\ & \quad - \beta r(c_{j,k}^*) \phi_{j,k} n_{j,k}^{\ell+1} - \frac{1}{\varepsilon^3} (1 - \phi_{j,k})(c_{j,k}^{\ell+1} - 1). \end{aligned}$$

4.3. Numerical boundary conditions

The boundary conditions on $\partial\tilde{\Omega}_{\text{bot}}$ are given by (3.6), which is implemented using $m = \phi n$ and the ghost cell technique as follows:

$$\overline{m}_{j,0}^{\ell} = \frac{\phi_{j,0}}{\phi_{j,1}} \overline{m}_{j,1}^{\ell}, \quad u_{j,0}^{\ell} = v_{j,0}^{\ell} = 0, \quad c_{j,0}^{\ell} = c_{j,1}^{\ell}, \quad \forall j, \ell.$$

The boundary conditions on $\tilde{\Gamma}$ are given by (3.5). We first rewrite the second equation in (3.5) as $\partial(\ln n)/\partial\boldsymbol{\nu} = \alpha\partial c/\partial\boldsymbol{\nu}$, which can be easily integrated on each of the three sides of $\tilde{\Gamma}$. We then use $m = \phi n$ and the ghost cell technique to end up with the following boundary conditions for the corresponding three sides:

$$\begin{aligned} \overline{m}_{0,k}^{\ell} & = \frac{\phi_{0,k}}{\phi_{1,k}} \overline{m}_{1,k}^{\ell} e^{\alpha(1-c_{1,k})}, \quad u_{0,k}^{\ell} = u_{1,k}^{\ell}, \quad v_{0,k}^{\ell} = v_{1,k}^{\ell}, \quad c_{0,k}^{\ell} = 1, \\ \overline{m}_{N+1,k}^{\ell} & = \frac{\phi_{N+1,k}}{\phi_{N,k}} \overline{m}_{N,k}^{\ell} e^{\alpha(1-c_{N,k})}, \quad u_{N+1,k}^{\ell} = u_{N,k}^{\ell}, \\ v_{N+1,k}^{\ell} & = v_{N,k}^{\ell}, \quad c_{N+1,k}^{\ell} = 1, \\ \overline{m}_{j,M+1}^{\ell} & = \frac{\phi_{j,M+1}}{\phi_{j,M}} \overline{m}_{j,M}^{\ell} e^{\alpha(1-c_{j,M})}, \quad u_{j,M+1}^{\ell} = u_{j,M}^{\ell}, \\ v_{j,M+1}^{\ell} & = v_{j,M}^{\ell}, \quad c_{j,M+1}^{\ell} = 1, \end{aligned}$$

for all j, k , and ℓ .

4.4. Summary of the numerical algorithm

Before demonstrating the performance of the proposed diffuse-domain-based numerical method, we summarize our approach for solving systems of PDEs in complicated domains in the following algorithm.

Algorithm 4.1. Hybrid finite-volume finite-difference method.

Data: The studied system of PDEs in a complex domain.

Step 1: Use the diffuse-domain method to reformulate this system properly.

Step 2: Perform asymptotic analysis of the reformulated system:

if the diffuse-domain model asymptotically converges to the original system

then

 | go to Step 3;

else

 | go back to Step 1;

end

Step 3: Numerically solve the modified system on a uniform Cartesian mesh in a large rectangle containing the original domain.

Result: Numerical solution of the studied system of PDEs.

5. Numerical Examples

In this section, we apply our new high-resolution method to simulate the bioconvection patterns of the oxygen-driven swimming bacteria in different sessile drops. In all of the examples, we use a uniform mesh with $\Delta x = \Delta y = 0.01$ and $\Delta t = 6.25 \times 10^{-6}$, which is chosen according to (4.19) and Remark 4.1. We fix the thickness of the diffuse-domain boundary to be $\varepsilon = 0.01$. We follow Ref. 44 and choose the cut-off function $r(c)$ being (1.2) with $c^* = 0.3$, and the following parameters: $\alpha = 10$, $\delta = 5$, and $Sc = 500$. The values of β and γ will vary and will be specified below.

The selection of the numerical experiments is motivated by the bacterial swimming and oxygen transport dynamics in sessile drops; see Ref. 44. We begin with Examples 1 and 2, where one can observe the formation of the stable stationary plumes (also shown in Ref. 15) in sessile drops of two different shapes as well as the behavior of the bacteria and fluid near solid-air-water contact lines. It is worth noting that the geometry of sessile drops is so complex that it is difficult to use standard numerical methods directly for such domains. Moreover, the plots of time evolution of kinetic energy are shown to verify the asymptotic stability of the system. We then increase the length of the drops in Examples 3 and 4 to study the influence of droplet geometry on the nonlinear dynamics of the model (1.1). In addition, we consider the case of larger cell density, which may result in more complicated bacteria propagation dynamics: Indeed, mushroom-shaped plumes will form in Examples 5 and 6, and this phenomenon is similar to the experimental one shown in Ref. 25. Finally, in Examples 7 and 8, we consider the cases of sessile drops surrounded by oxygen (this would occur when droplets are placed on superhydrophobic surfaces): Bioconvection phenomena in such cases are also interesting to study.

5.1. Stable stationary plumes

In this section, we consider four sessile drops of different shapes determined by a given function $f(x, y)$ representing the original domain $\Omega = \{(x, y) \mid f(x, y) > 0, y > 0\}$, for which we compute the signed distance function $d(x, y)$ to $\Gamma = \{(x, y) \mid f(x, y) = 0, y > 0\}$ needed to obtain the diffuse-domain function $\phi(x, y)$; see (3.7). In order to implement the proposed diffuse-domain-based method, Ω is imbedded into a larger domain $\tilde{\Omega}$, which is taken either $\tilde{\Omega} = [-5, 5] \times [0, 1.5]$ (Examples 1 and 2) or $\tilde{\Omega} = [-7.5, 7.5] \times [0, 1.5]$ (Examples 3 and 4).

In Examples 1–4, we take the parameters $\beta = 10$ and $\gamma = 1000$.

Example 1. In the first example, we solve the system (3.1)–(3.8) subject to the following initial data:

$$n(x, y, 0) = \begin{cases} 1 & \text{if } y > 0.499 - 0.01 \sin(\pi(x - 1.5)), \\ 0.5 & \text{otherwise,} \end{cases}$$

$$c(x, y, 0) \equiv 1, \quad u(x, y, 0) = v(x, y, 0) \equiv 0,$$

which is prescribed in the domain Ω determined by

$$f(x, y) = \begin{cases} 4.8 + x - (0.9y + 0.2)^2 - 0.1(0.9y + 0.2)^{16} & \text{if } x \leq 0, \\ 4.8 - x - (0.9y + 0.2)^2 - 0.1(0.9y + 0.2)^{16} & \text{otherwise;} \end{cases}$$

see the upper left panel in Fig. 3, where the shape of the drop and initial cell density are plotted.

The time evolution of the cell density n is also shown in Fig. 3. As one can see, the bacteria first ($t = 0.1$) aggregate along the boundary Γ as the concentration




Fig. 3. (Color online) Example 1: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.




Fig. 4. (Color online) Example 1: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.

of the oxygen is high there, but then ($t = 0.2$) the gravity forces start playing an essential role and some of the bacteria fall down forming the plumes ($t = 0.3$) both at the corners and in the middle of the drop. Later on, the shape of the plumes slightly changes ($t = 1$) and by time $t = 2$ the plumes are already stationary (compare with the cell density at the very large time $t = 6$). For the sake of brevity, we plot the c -component of the computed solution only at time $t = 6$. The obtained results confirm the ability of the proposed numerical method to capture stationary plumes in a stable manner.

In order to numerically verify the stability of the plumes, we check the time evolution of the kinetic energy (see Fig. 4), which clearly converges to a constant value. In addition, we plot the velocity field together with the $n = 0.7$ cell density level set at time $t = 1$ (see Fig. 5), which illustrate how the stationary plumes are supported by the fluid.

We note that the obtained results are in a very good qualitative agreement with the results reported in Ref. 15, where the system (1.1) was considered in a rectangular domain subject to the periodic boundary conditions in the horizontal direction. In addition, the diffuse-domain-based numerical method proposed here is capable of treating non-rectangular domains and resolving the accumulation layers at the drop corners and creation of vortices there.




Fig. 5. (Color online) Example 1: Velocity field \mathbf{u} and the $n = 0.7$ level set at time $t = 1$.




Fig. 6. (Color online) Example 2: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.

Example 2. Next, we consider the same initial setting as in Example 1 but in a sessile drop of a different shape determined by

$$f(x, y) = \begin{cases} 4.8 + x - |1.5y - 0.75|^{2.5} - (1.5y - 0.75)^{10} & \text{if } x \leq 0, \\ 4.8 - x - |1.5y - 0.75|^{2.5} - (1.5y - 0.75)^{10} & \text{otherwise.} \end{cases}$$

Compared with the drop in the previous example, this one has rounded edges while still having a flat bottom interface; see the upper left panel in Fig. 6, where the shape of the drop and initial cell density are plotted.

The time evolution of the cell density n is also shown in Fig. 6. As one can see, the evolution process is similar to the one in Example 1 with the only exception that the structure of the aggregated area at the edges of the drop is different as the oxygen supply is available underneath that part of the drop considered here. The solution converges to a stationary state containing bacteria plumes, which can be seen in the bottom row of Fig. 6 (the oxygen concentration c at the final time $t = 6$ is also shown there). The convergence towards the steady state is also confirmed by following the time evolution of the kinetic energy (Fig. 7), which clearly flattens by time $t = 1$. As in Example 1, we also plot the velocity field together with the $n = 0.7$ cell density level set at time $t = 1$ (Fig. 8), which illustrate how the stationary plumes are supported by the fluid.

Once again, we emphasize that the proposed diffuse-domain-based numerical method is capable of numerically solving the fluid-chemotaxis system in rather complicated domains.




Fig. 7. (Color online) Example 2: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.




Fig. 8. (Color online) Example 2: Velocity field \mathbf{u} and the $n = 0.7$ level set at time $t = 1$.

Example 3. In the third example, we consider the same initial setting as in Example 1 but the drop is now longer. Its precise shape is determined by

$$f(x, y) = \begin{cases} 4.8 + \frac{2}{3}x - (0.9y + 0.2)^2 - 0.1(0.9y + 0.2)^{16} & \text{if } x \leq 0, \\ 4.8 - \frac{2}{3}x - (0.9y + 0.2)^2 - 0.1(0.9y + 0.2)^{16} & \text{otherwise;} \end{cases}$$

see the upper left panel in Fig. 9, where the shape of the drop and initial cell density are plotted. The time evolution of the cell density n as well as the profile of the oxygen concentration c at the final time $t = 2$, by which the solution reaches its steady state, are also shown in Fig. 9. As one can see, the proposed numerical method can handle longer drops and the only qualitative difference between the steady states here and in Example 1 is in the number of plumes emerging during the evolution process.

Example 4. The final example of this section is a modification of Example 2 as we now take a longer drop determined by

$$f(x, y) = \begin{cases} 4.8 + \frac{2}{3}x - |1.5y - 0.75|^{2.5} - (1.5y - 0.75)^{10} & \text{if } x \leq 0, \\ 4.8 - \frac{2}{3}x - |1.5y - 0.75|^{2.5} - (1.5y - 0.75)^{10} & \text{otherwise;} \end{cases}$$




Fig. 9. (Color online) Example 3: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.

see the upper left panel in Fig. 10, where the shape of the drop and initial cell density are plotted. The time evolution of n , which converges to the steady state by $t = 2$, together with the profile of c at the final time can be also seen in Fig. 10. The obtained stationary solution contains two additional plumes compared with the solution reported in Example 2, but rather than this these two solutions are qualitatively similar, which confirms the robustness of our numerical method.

5.2. Mushroom-shaped plumes for high-density data

In this section, we choose the parameters $\beta = 100$ and $\gamma = 10,000$, which correspond to a 10-times larger reference cell density n_r ; see (2.2). The evolution process will now be substantially faster so that we conduct the simulations for a shorter period and take the final time $t = 0.5$.




Fig. 10. (Color online) Example 4: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.

The goal of the simulations reported in Examples 5 and 6 is to demonstrate the ability of the proposed diffuse-domain-based numerical method to handle more complicated bacteria propagation dynamics, which are expected to occur when the reference cell density n_r is larger.

Example 5. In this example, we use precisely the same shape of the drop and initial data as in Example 1. The time evolution of the computed cell density n is shown in Fig. 11. As one can see, compared with Example 1 heavier mushroom-shaped plumes are formed by time $t = 0.08$. Later on (by time $t = 0.1$) these plumes are disintegrated and a part of the bacteria fall to the bottom of the drop and become inactive due to the low oxygen concentration there. After that, smaller mushroom-shaped plumes are re-emerged and then disintegrate several times. At the same time, one can observe the propagation of the bacteria along the top part of the drop towards its corners. Eventually, the evolution process seems to converge to the steady state by the final time $t = 0.5$ as confirmed by the stabilization of the




Fig. 11. (Color online) Example 5: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.




Fig. 12. (Color online) Example 5: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.

kinetic energy by then; see Fig. 12. It may also be instructive to see the final time distribution of the oxygen concentration c (see the bottom right panel in Fig. 11), which indicates that after falling down the bacteria in the lower part of the drop remain inactive.




Fig. 13. (Color online) Example 6: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.

Example 6. In the next example, we use precisely the same shape of the drop and initial data as in Example 2. Time snapshots of the computed cell density n at different times are shown in Fig. 13. In principle, the time evolution is quite similar to what was observed in Example 5, but due to the difference in the shape of the drops, several distinctive features can be seen. While the mushroom-type




Fig. 14. (Color online) Example 6: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.

plumes formed at about $t = 0.08$ are qualitatively similar to those in Fig. 11, the solution at later time develops a different symmetry: by the time $t = 0.16$ – 0.17 three plumes (one in the center of the drop and two plumes propagating to the sides) are formed. Later on they keep disintegrating and re-appearing and by $t = 0.24$, one can see only one plume, which remained in the center of the drop as the other two plumes practically merged with the top boundary cell layer. After that, the remaining plume keeps disintegrating and re-emerging until the solution reaches its steady state. This convergence is confirmed by the stabilization of the kinetic energy (see Fig. 14) and also by the final time oxygen distribution (see the bottom right panel in Fig. 13).

5.3. Plumes in sessile drops surrounded by oxygen

In this section, we consider the sessile drop surrounded by oxygen. The shape of the drop is determined by the function

$$f(x, y) = \begin{cases} 4.8 + x - |1.5y - 0.95|^{2.5} - (1.5y - 0.95)^{10} & \text{if } x \leq 0, \\ 4.8 - x - |1.5y - 0.95|^{2.5} - (1.5y - 0.95)^{10} & \text{otherwise,} \end{cases}$$

representing the original domain $\Omega = \{(x, y) \mid f(x, y) > 0, y > 0.1\}$, for which we compute the signed distance function $d(x, y)$ to

$$\partial\Omega = \{(x, y) \mid f(x, y) = 0, y > 0.1\} \cup \{(x, y) \mid f(x, y) > 0, y = 0.1\}$$

needed to obtain the diffuse-domain function $\phi(x, y)$ in (3.7). In order to implement the proposed diffuse-domain-based method, Ω is imbedded into $\tilde{\Omega} = [-5, 5] \times [0, 1.5]$.

Unlike the drops considered up to now, here we model the drop surrounded by oxygen. Therefore, the boundary conditions

$$\boldsymbol{\nu} \cdot \mathbf{u} = 0, \quad \boldsymbol{\nu} \cdot \nabla(\mathbf{u} \cdot \boldsymbol{\tau}) = 0, \quad (\alpha n \nabla c - \nabla n) \cdot \boldsymbol{\nu} = 0, \quad c = 1, \quad \forall (x, y) \in \partial\Omega,$$

which were used along the top portion Γ in (2.8), are now set along the entire boundary $\partial\Omega$.

We solve the system (3.1)–(3.4), (3.7), (3.8) subject to the boundary conditions

$$\boldsymbol{\nu} \cdot \nabla \mathbf{u} = \mathbf{0}, \quad (\alpha n \nabla c - \nabla n) \cdot \boldsymbol{\nu} = 0, \quad c = 1, \quad \forall (x, y) \in \partial\tilde{\Omega}$$




Fig. 15. (Color online) Example 7: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.

instead of the previously used (3.5)–(3.6), and the following initial data:

$$n(x, y, 0) = \begin{cases} 1 & \text{if } y > 0.599 - 0.01 \sin(\pi(x - 1.5)), \\ 0.5 & \text{otherwise,} \end{cases}$$

$$c(x, y, 0) \equiv 1, \quad u(x, y, 0) = v(x, y, 0) \equiv 0.$$

Example 7. In this example, we take $\beta = 20$ and $\gamma = 2000$ and we compute the solutions until the final time $t = 5$. In Fig. 15, the computed cell densities at different times are plotted along with the oxygen concentration, which is shown at the final time only. The major difference between this example and Examples 1–6 is that the oxygen is now accessible around the entire boundary of the drop. Therefore, the bacteria immediately start propagating along the boundary towards the lower part of the drop (this can be seen even at a small time $t = 0.1$). At the same time, the gravity causes the formation of the plumes (see, e.g. the solution at $t = 0.2$). These plumes are unstable and later on more plumes are formed. At larger times, a small plume at the center of the drop is merged and it seems to be stable as the solution converges to its steady state by the final time; see also Fig. (16), where the kinetic energy is depicted.

Example 8. The final example is similar to Example 7 with the only exception that here we take $\beta = 40$ and $\gamma = 4000$. These values correspond to a twice larger reference cell density n_r , which leads to a faster dynamics. Indeed, as one can clearly see




Fig. 16. (Color online) Example 7: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.




Fig. 17. (Color online) Example 8: Time snapshots of the computed cell densities n at different times and the computed oxygen concentration c at the final time.




Fig. 18. (Color online) Example 8: Time-evolution of the kinetic energy $\|\mathbf{u}\|_{L^2(\Omega)}$.

from Figs. 17 and 18, the solution converges to its steady state substantially faster than in Example 7. It should also be observed that the obtained steady state qualitatively different from the one reported in Fig. 15: the steady state now contains two plumes (not only one plume as in the previous example) and there is a slightly larger concentration of bacteria in the internal part of the drop (this can be clearly seen from the final time oxygen distribution shown in the lower right panel of Fig. 17).

6. Conclusion

In this paper, we have introduced a new positivity preserving and high-resolution method for the coupled chemotaxis-fluid system in a sessile drop. Our method is based on the diffuse-domain approach, which is implemented to derive a cf-DD model. We have shown that the obtained cf-DD system converges to the original chemotaxis-fluid system as the thickness of the diffuse-domain interface shrinks to zero. In order to numerically solve the resulting cf-DD system, we have developed a second-order hybrid finite-volume finite-difference method, which preserves non-negativity of the computed cell density.

We have tested the proposed diffuse-domain-based method on a number of numerical experiments, in which we have not only demonstrated the ability of our method to handle complex computational domains, but also systematically studied bacteria collective behavior in sessile droplets of a variety of different shapes. It has been observed that when the amount of bacteria is moderate, stable stationary plumes are formed inside the droplet and substantial amount of the bacteria will aggregate in the corners of the droplet while creating vortices there. When the amount of bacteria is increased, the finger-like plumes flare out into mushroom-shaped plumes, which are, however, unstable and disintegrate in time. At the same time, the solutions converge to nontrivial steady states in all of the studied examples. Based on the obtained numerical results, we conjecture that the evolution of bacteria is related to both the total amount of bacteria in the droplet and the shape of the droplet. The main goal of the presented simulations is to demonstrate that the proposed numerical method can provide one with a valuable insight on the bacteria collective behavior in complex geometries, whose detailed study is left for the future work.

Acknowledgments

The work of A. Chertock was supported in part by the NSF Grants DMS-1818684 and DMS-2208438. The work of A. Kurganov was supported in part by the NSFC Grants 12111530004 and 12171226, and by the fund of the Guangdong Provincial Key Laboratory of Computational Science and Material Design (No. 2019B030301001). The work of Zhen Zhang was partially supported by the NSFC Grant 12071207, NSFC Tianyuan-Pazhou Grant 12126602, the Natural Science Foundation of Guangdong Province (No. 2021A1515010359), and the Guangdong Provincial Key Laboratory of Computational Science and Material Design (No. 2019B030301001).

References

1. H. Abels, H. Garcke and G. Grün, Thermodynamically consistent, frame indifferent diffuse interface models for incompressible two-phase flows with different densities, *Math. Models Methods Appl. Sci.* **22** (2012) 1150013.

2. S. Aland, J. Lowengrub and A. Voigt, Two-phase flow in complex geometries: A diffuse domain approach, *CMES Comput. Model. Eng. Sci.* **57** (2010) 77–107.
3. S. Aland, J. Lowengrub and A. Voigt, A continuum model of colloid-stabilized interfaces, *Phys. Fluids* **23** (2011) 062103.
4. D. M. Anderson, G. B. McFadden and A. A. Wheeler, A phase-field model with convection: Sharp-interface asymptotics, *Phys. D* **151** (2001) 305–331.
5. G. Arumugam and J. Tyagi, Keller–Segel chemotaxis models: A review, *Acta Appl. Math.* **171** (2021) 6.
6. N. Bellomo, A. Bellouquid and N. Chouhad, From a multiscale derivation of nonlinear cross-diffusion models to Keller–Segel models in a Navier–Stokes fluid, *Math. Models Methods Appl. Sci.* **26** (2016) 2041–2069.
7. N. Bellomo, L. Gibelli and N. Outada, On the interplay between behavioral dynamics and social interactions in human crowds, *Kinet. Relat. Models* **12** (2019) 397–409.
8. N. Bellomo, N. Outada, J. Soler, Y. Tao and M. Winkler, Chemotaxis and cross-diffusion models in complex environments: Models and analytic problems toward a multiscale vision, *Math. Models Methods Appl. Sci.* **32** (2022) 713–792.
9. N. Bellomo, K. J. Painter, Y. Tao and M. Winkler, Occurrence versus absence of taxis-driven instabilities in a May–Nowak model for virus infection, *SIAM J. Appl. Math.* **79** (2019) 1990–2010.
10. C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers. I. Asymptotic Methods and Perturbation Theory* (Springer-Verlag, 1999) Reprint of the 1978 original.
11. M. Braukhoff and B. Q. Tang, Global solutions for chemotaxis-Navier–Stokes system with Robin boundary conditions, *J. Differential Equations* **269** (2020) 10630–10669.
12. A. Bueno-Orovio and V. M. Pérez-García, Spectral smoothed boundary methods: The role of external boundary conditions, *Numer. Methods Partial Differential Equations* **22** (2006) 435–448.
13. A. Bueno-Orovio, V. M. Pérez-García and F. H. Fenton, Spectral methods for partial differential equations in irregular domains: The spectral smoothed boundary method, *SIAM J. Sci. Comput.* **28** (2006) 886–900.
14. S. Chakraborty, F. Ivančić, M. Solovchuk and T. W. H. Sheu, Stability and dynamics of a chemotaxis system with deformed free-surface in a shallow chamber, *Phys. Fluids* **30** (2018) 071904.
15. A. Chertock, K. Fellner, A. Kurganov, A. Lorz and P. A. Markowich, Sinking, merging and stationary plumes in a coupled chemotaxis-fluid model: A high-resolution numerical approach, *J. Fluid Mech.* **694** (2012) 155–190.
16. Y. Deleuze, C.-Y. Chiang, M. Thiriet and T. W. H. Sheu, Numerical study of plume patterns in a chemotaxis–diffusion–convection coupling system, *Comput. Fluids* **126** (2016) 58–70.
17. C. Dombrowski, L. Cisneros, S. Chatkaew, R. E. Goldstein and J. O. Kessler, Self-concentration and large-scale coherence in bacterial dynamics, *Phys. Rev. Lett.* **93** (2004) 098103.
18. A. Duarte-Rodríguez, M. A. Rodríguez-Bellido, D. A. Rueda-Gómez and E. J. Villamizar-Roa, Numerical analysis for a chemotaxis-Navier–Stokes system, *ESAIM Math. Model. Numer. Anal.* **55** (2021) S417–S445.
19. S. Gottlieb, C.-W. Shu and E. Tadmor, Strong stability-preserving high-order time discretization methods, *SIAM Rev.* **43** (2001) 89–112.
20. J. L. Guermond and J. Shen, On the error estimates for the rotational pressure-correction projection methods, *Math. Comp.* **73** (2004) 1719–1737.

21. Z. Guo, P. Lin, J. Lowengrub and S. M. Wise, Mass conservative and energy stable finite difference methods for the quasi-incompressible Navier–Stokes–Cahn–Hilliard system: Primitive variable and projection-type schemes, *Comput. Methods Appl. Mech. Engrg.* **326** (2017) 144–174.
22. Z. Guo, F. Yu, P. Lin, S. Wise and J. Lowengrub, A diffuse domain method for two-phase flows with large density ratio in complex geometries, *J. Fluid Mech.* **907** (2021) A38.
23. N. A. Hill and T. J. Pedley, Bioconvection, *Fluid Dynam. Res.* **37** (2005) 1–20.
24. T. Hillen and K. J. Painter, A user’s guide to PDE models for chemotaxis, *J. Math. Biol.* **58** (2009) 183–217.
25. A. J. Hillesdon, T. J. Pedley and J. O. Kessler, The development of concentration gradients in a suspension of chemotactic bacteria, *Bull. Math. Biol.* **57** (1995) 299–344.
26. M. H. Holmes, *Introduction to Perturbation Methods*, Texts in Applied Mathematics, 2nd edn., Vol. 20 (Springer, 2013).
27. D. Horstmann, From 1970 until present: The Keller–Segel model in chemotaxis and its consequences. I, *Jahresber. Deutsch. Math.-Verein.* **105** (2003) 103–165.
28. X. Huang, X. Feng, X. Xiao and K. Wang, Fully decoupled, linear and positivity-preserving scheme for the chemotaxis-Stokes equations, *Comput. Methods Appl. Mech. Engrg.* **383** (2021) 113909.
29. F. Ivančić, T. W. H. Sheu and M. Solovchuk, The free surface effect on a chemotaxis-diffusion-convection coupling system, *Comput. Methods Appl. Mech. Engrg.* **356** (2019) 387–406.
30. F. Ivančić, T. W. H. Sheu and M. Solovchuk, Bacterial chemotaxis in thin fluid layers with free surface, *Phys. Fluids* **32** (2020) 061902.
31. Y. Ke, J. Li and Y. Wang, *Analysis of Reaction-Diffusion Models with the Taxis Mechanism*, Financial Mathematics and Fintech (Springer, 2022).
32. E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an instability, *J. Theoret. Biol.* **26** (1970) 399–415.
33. E. F. Keller and L. A. Segel, Model for chemotaxis, *J. Theoret. Biol.* **30** (1971) 225–234.
34. K. Y. Lervåg and J. Lowengrub, Analysis of the diffuse-domain method for solving PDEs in complex geometries, *Commun. Math. Sci.* **13** (2015) 1473–1500.
35. X. Li, J. Lowengrub, A. Rätz and A. Voigt, Solving PDEs in complex geometries: A diffuse domain approach, *Commun. Math. Sci.* **7** (2009) 81–107.
36. A. Lorz, A coupled Keller–Segel–Stokes model: Global existence for small initial data and blow-up delay, *Commun. Math. Sci.* **10** (2012) 555–574.
37. H. Nessyahu and E. Tadmor, Nonoscillatory central differencing for hyperbolic conservation laws, *J. Comput. Phys.* **87** (1990) 408–463.
38. C. S. Patlak, Random walk with persistence and external bias, *Bull. Math. Biophys.* **15** (1953) 311–338.
39. R. L. Pego, Front migration in the nonlinear Cahn–Hilliard equation, *Proc. R. Soc. Lond. Ser. A* **422** (1989) 261–278.
40. J. Shen, Modeling and numerical approximation of two-phase incompressible flows by a phase-field approach, in *Multiscale Modeling and Analysis for Materials Simulation*, Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, Vol. 22 (World Scientific Publishing, 2012), pp. 147–195.
41. P. K. Sweby, High resolution schemes using flux limiters for hyperbolic conservation laws, *SIAM J. Numer. Anal.* **21** (1984) 995–1011.

42. K. E. Teigen, X. Li, J. Lowengrub, F. Wang and A. Voigt, A diffuse-interface approach for modeling transport, diffusion and adsorption/desorption of material quantities on a deformable interface, *Commun. Math. Sci.* **7** (2009) 1009–1037.
43. K. E. Teigen, P. Song, J. Lowengrub and A. Voigt, A diffuse-interface method for two-phase flows with soluble surfactants, *J. Comput. Phys.* **230** (2011) 375–393.
44. I. Tuval, L. Cisneros, C. Dombrowski, C. W. Wolgemuth, J. O. Kessler and R. E. Goldstein, Bacterial swimming and oxygen transport near contact lines, *Proc. Natl. Acad. Sci. USA* **102** (2005) 2277–2282.
45. B. van Leer, Towards the ultimate conservative difference scheme. V. A second-order sequel to Godunov’s method, *J. Comput. Phys.* **32** (1979) 101–136.
46. M. Winkler, How far do chemotaxis-driven forces influence regularity in the Navier–Stokes system? *Trans. Amer. Math. Soc.* **369** (2017) 3067–3125.
47. M. Winkler, A three-dimensional Keller–Segel–Navier–Stokes system with logistic source: Global weak solutions and asymptotic stabilization, *J. Funct. Anal.* **276** (2019) 1339–1401.
48. J.-J. Xu, W. Shi and M.-C. Lai, A level-set method for two-phase flows with soluble surfactant, *J. Comput. Phys.* **353** (2018) 336–355.
49. F. Yu, Z. Guo and J. Lowengrub, Higher-order accurate diffuse-domain methods for partial differential equations with Dirichlet boundary conditions in complex, evolving geometries, *J. Comput. Phys.* **406** (2020) 109174, 34.
50. H.-C. Yu, H.-Y. Chen and K. Thornton, Extended smoothed boundary method for solving partial differential equations with general boundary conditions on complex boundaries, *Model. Simul. Mater. Sci.* **20** (2012) 075008.